

lighten your day

the Velphoro way

Velphoro, **the strongest phosphate binder**, helps you manage your phosphorus levels—more control, fewer pills, more ways to take them.



INDICATION

VELPHORD

Velphoro[®] (sucroferric oxyhydroxide) is a phosphate binder indicated for the control of serum phosphorus levels in adult and pediatric patients 9 years of age and older with chronic kidney disease on dialysis.



(sucroferric oxyhydroxide) chewable tablets

Phosphorus is a mineral found in your body and in foods you eat.

When you have chronic kidney disease and are on dialysis, your body cannot get rid of excess phosphorus.

Diet and dialysis are two ways to keep phosphorus from building up in your body. When diet and dialysis aren't enough, your doctor may prescribe a phosphate-lowering medicine like Velphoro.

Look for the letters "phos." There is hidden phosphorus (or phosphate) in many foods.

INGREDIENTS: POTATOES, SUNFLOWER OIL, AND/OR CORN OIL, SALT, SODIUM DIACETATE, CHEESE CULTURES, ENZYMES, BUTTER, MILK POWDER, SUGAR, SOYBEAN OIL, WHEY, CITRIC ACID, DISODIUM PHOSPHATE, OLIVE OIL.

Why your doctor has prescribed Velphoro

Velphoro is a phosphate binder, which helps people with chronic kidney disease on dialysis control the level of phosphorus in their blood.

Velphoro only works if you take it with food.

Velphoro works **like a magnet to pull the phosphorus out of the food** you've eaten and help remove it from your body.

Please see Important Safety Information on page 12, and full Prescribing Information enclosed in pocket.



(sucroferric oxyhydroxide) chewable tablets

Velphoro is the strongest phosphate binder, so you can take fewer pills. In fact, one tablet of Velphoro locks up about twice as much phosphorus as other commonly used binders.



Because Velphoro is so strong, most people take only 3 or 4 Velphoro tablets per day compared with 9-12 tablets per day with other phosphate binders.



Velphoro helps more people get to and stay at goal

Real-world data show that when people switched to Velphoro from another phosphate binder, double the percentage of people were able to control their phosphorus levels with **half** the pills.



Real-world data from people who stayed on Velphoro for 2 years showed that 40% more of them stayed at their phosphorus goals compared with people who switched to another phosphate binder.



Taking Velphoro as prescribed may help you get to your phosphorus goal and stay there.

Please see Important Safety Information on page 12, and full Prescribing Information enclosed in pocket.

VELPHORO[®] (sucroferric oxyhydroxide) chewable tablets

Take Velphoro when you eat



Take Velphoro when you eat a meal or snack, as prescribed by your doctor. You will probably start on Velphoro with 1 pill every time you eat a meal.

Your doctor may adjust your dose after a while, based on how your phosphate levels are tracking.

After a year or two, most people still only take about **4 Velphoro tablets per day**, **not per meal**, like with some other phosphate binders.

You choose how to take Velphoro

Velphoro lightens up mealtime because you can take it different ways. It's easy to break, chew, or crush, so it's not obvious to others that you're taking medicine.

Velphoro is not meant to be swallowed whole. Just "break and take." Take it just before, during, or just after a meal or snack.



What if you forget?

If you forget to take Velphoro, take your regular dose with your next meal. Remember, it only works if you take it with food!



sucroferric oxyhydroxide) chewable tablets

Side effects with Velphoro are mostly mild and go away on their own after a short time.

Most common is diarrhea, which usually resolves by itself. Make sure that you tell your healthcare team if you're currently taking a laxative or stool softener. Some people had nausea. Because Velphoro is made from iron, it may turn your stools black. This is normal—and expected—just as it is with other medicines that contain iron.

Some people have reported discoloration of their teeth, or a rash.

Most people reported no side effects with Velphoro, but some did. Very few people have stopped taking Velphoro because of side effects.

Please see Important Safety Information on page 12, and full Prescribing Information enclosed in pocket.

Speak to your healthcare team if side effects become a concern.

VELPHORD



How to remember to take Velphoro

Meals Or snacks Velphoro Every time*

Use these tips



FORWARD

- Keep Velphoro near where you eat⁺
- Put a note on your refrigerator
- Set an alarm on your phone to ring 3 times a day at mealtimes
- Mark your calendar to refill Velphoro a week before you run out

Ask your healthcare team for their favorite tips

*Take Velphoro when you eat a meal or snack as prescribed by your doctor. *Always keep the bottle or pill container of Velphoro tightly closed. Store at room temperature, and below 87° F.

Visit the Velphoro Video Center

Hear people talk about their personal dialysis journeys and how they succeeded with Velphoro.



"I would recommend Velphoro because you can break it, crush it, chew it, and it tastes great." —Anita



"I could take one phosphate binder instead of six with each meal."

-Antonio



"Velphoro helps me control my phosphorus and maintain it." —Easter



"Ever since we switched my phosphate binder, my phosphorus has been spot on." -Robert



Watch their stories at VelphoroVoices.com.

Scan the code with your cell phone camera

Please see Important Safety Information on page 12, and full Prescribing Information enclosed in pocket.

The paid testimonials provided are representative of that individual's experience, and not all patients may experience these benefits.

IMPORTANT SAFETY INFORMATION

Velphoro chewable tablets must be taken with meals. Velphoro should be chewed or crushed. Do not swallow whole. Tell your healthcare provider about all the medicines you take, including prescription and nonprescription medicines, vitamins, and other supplements. Velphoro can interact with other medicines.

Tell your healthcare provider if you have any of the following: peritonitis (an infection) during peritoneal dialysis, significant gastric or liver disorder, recent major gastrointestinal (GI) surgery, a history of hemochromatosis or other disease that results in iron buildup in the body. People with these conditions were not included in clinical studies with Velphoro, and your healthcare provider will monitor your iron levels while you are taking Velphoro.

Velphoro can cause side effects. The most common side effects are discolored stool, diarrhea, and nausea. Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

Discolored (black) stool is considered normal if you are taking an oral medication that contains iron, like Velphoro. Discolored stool may mask gastrointestinal bleeding, which was not a side effect of Velphoro in clinical studies.

Before taking Velphoro, tell your doctor if you are pregnant, plan to become pregnant, or are breastfeeding.

Velphoro is available by prescription only. For additional Safety Information, please see full Prescribing Information.

To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Medical Care Customer Service at 1-800-323-5188 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.





Velphoro has been used for more than 10 years with thousands of people on dialysis like you.



lighten your day with Velphoro

- Velphoro is the strongest phosphate binder, so you can take fewer pills per meal
- Velphoro helps more people get to goal with fewer pills
- You can choose how to take Velphoro. You can break it, chew it, or crush it
- Take Velphoro as prescribed and make it part of your daily meal plan

Talk to your healthcare team for more information about how Velphoro can help you control your phosphorus. For more information visit Velphoro.com.



RENAL PHARMACEUTICALS

© 2018, 2025 Fresenius Medical Care. All Rights Reserved. Fresenius Medical Care and the triangle logo are trademarks for which Fresenius Medical Care AG holds rights. Velphoro is a trademark of Vifor Fresenius Medical Care Renal Pharma Ltd. All other trademarks are the property of their respective owners.

Distributed by: Fresenius Medical Care North America, 920 Winter Street, Waltham, MA 02451-1457

US-Velphoro-000166 Rev B 01/2025

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use VELPHORO® safely and effectively. See full prescribing information for VELPHOR0[®].

VELPHOR0® (sucroferric oxyhydroxide) chewable tablet for oral use Initial U.S. Approval: 2013

RECENT MAJOR CHANGES		
Indications and Usage (1)	7/2024	
Dosage and Administration (2.1)	7/2024	

----- INDICATIONS AND USAGE ------Velphoro is a phosphate binder indicated for the control of serum phosphorus levels in adult and pediatric patients 9 years of age and older with chronic kidney disease on dialysis. (1)

- ----- DOSAGE AND ADMINISTRATION ------
- Chew or crush Velphoro tablets, do not swallow whole. (2.2)
- The recommended starting dose for adults and pediatric patients 12 years of age and older is one 500 mg tablet three times daily with meals. (2.1)
- The recommended starting dose for pediatric patients 9 to <12 years of age is one 500 mg tablet two times daily with meals. (2.1)
- Adjust dosage by one 500 mg tablet per day as needed until an acceptable serum phosphorus level is reached, with regular monitoring afterwards. Titrate as often as weekly. (2.1)

----- DOSAGE FORMS AND STRENGTHS ------

Velphoro chewable tablet 500 mg (3)

FULL PRESCRIBING INFORMATION: CONTENTS*

INDICATIONS AND USAGE

- DOSAGE AND ADMINISTRATION 2
 - 2.1 General Dosing Information
 - 2.2 Administration
- **DOSAGE FORMS AND STRENGTHS** 3
- 4 CONTRAINDICATIONS
- WARNINGS AND PRECAUTIONS 5
- 5.1 Monitoring in Patients with Gastrointestinal Disorders or Iron Accumulation Disorders
- **ADVERSE REACTIONS**

6

- 6.1 Clinical Trial Experience
- 6.2 Postmarketing Experience
- 7 DRUG INTERACTIONS

8 **USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use

FULL PRESCRIBING INFORMATION

INDICATIONS AND USAGE

Velphoro is indicated for the control of serum phosphorus levels in adults and pediatric patients 9 years of age and older with chronic kidney disease (CKD) on dialysis.

DOSAGE AND ADMINISTRATION 2

General Dosing Information 2.1

The recommended starting dose of Velphoro in adults and pediatric patients 12 years of age and older is one 500 mg tablet three times daily with meals. The recommended starting dose of Velphoro in pediatric patients 9 to <12 years of age is one 500 mg tablet two times daily with meals.

Monitor serum phosphorus levels and titrate the dose of Velphoro in decrements or increments of one 500 mg tablet per day as needed until an acceptable serum phosphorus level is reached, with regular monitoring afterwards. Titrate as often as weekly. If the recommended daily dose cannot be divided equally among meals, administer the larger dose with the largest meal of the day.

In clinical studies, on average, adult patients and pediatric patients 12 years of age and older required 1,500 mg to 2,000 mg (3 to 4 tablets) a day to control serum phosphorus levels; pediatric patients 9 to <12 years of age required on average 1,500 mg (3 tablets) to control serum phosphorus levels. Daily doses as high as 3,000 mg per day have been studied in adults and pediatric patients 9 years of age and older.

2.2 Administration

Velphoro chewable tablets should be chewed or crushed. Do not swallow whole.

Velphoro chewable tablets must be administered with meals

If one or more doses of Velphoro are missed, the medication should be resumed with the next food intake.

DOSAGE FORMS AND STRENGTHS

Velphoro is supplied in a strength of 500 mg as a brown, circular, bi-planar chewable tablet embossed with "PA 500" on one side.

Each chewable tablet contains 500 mg iron (equivalent to 2,500 mg sucroferric oxyhydroxide).

----- CONTRAINDICATIONS ------

• None (4)

------ WARNINGS AND PRECAUTIONS ------

· Patients with peritonitis during peritoneal dialysis, significant gastric or hepatic disorders, history of recent major gastrointestinal surgery, or with a history of hemochromatosis or other diseases with iron accumulation have not been included in clinical studies with Velphoro. Monitor effect and iron homeostasis in such patients. (5.1)

----- ADVERSE REACTIONS ------

The most common adverse drug reactions to Velphoro chewable tablets in clinical trials were discolored feces (12%) and diarrhea (6%). (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Medical Care North America at 1-800-323-5188 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- DRUG INTERACTIONS ------

- · Velphoro can be administered concomitantly with oral calcitriol, ciprofloxacin, digoxin, enalapril, furosemide, HMG-CoA reductase inhibitors, hydrochlorothiazide, losartan, metoprolol, nifedipine, omeprazole, quinidine and warfarin. (7)
- Take acetylsalicylic acid, cephalexin and doxycycline at least 1 hour before Velphoro. (7)
- Take levothyroxine at least 4 hours before Velphoro. (7)
- For oral medications not listed above where a reduction of bioavailability would be ٠ clinically significant consider separation of the timing of administration. Consider monitoring clinical responses or blood levels of the concomitant medication (7)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 8/2024

- 10 OVERDOSAGE
- 11 DESCRIPTION
- **CLINICAL PHARMACOLOGY** 12
 - 12.1 Mechanism of Action
 - 12.2 Pharmacodynamics
 - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY
- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility 14
 - **CLINICAL STUDIES**
 - 14.1 Fixed-dose Study
 - 14.2 Dose Titration Study
 - 14.3 Pediatric Study
- HOW SUPPLIED/STORAGE AND HANDLING 16 16.1 How Supplied 16.2 Storage
- PATIENT COUNSELING INFORMATION 17

*Sections or subsections omitted from the full prescribing information are not listed.

CONTRAINDICATIONS 4

None.

WARNINGS AND PRECAUTIONS 5

5.1 Monitoring in Patients with Gastrointestinal Disorders or Iron Accumulation Disorders

Each chewable tablet contains 500 mg iron (equivalent to 2,500 mg sucroferric oxyhydroxide) [see Clinical Pharmacology (12.3)]. Patients with peritonitis during peritoneal dialysis, significant gastric or hepatic disorders, following major gastrointestinal (GI) surgery, or with a history of hemochromatosis or other diseases with iron accumulation have not been included in clinical studies with Velphoro.

Monitor effect and iron homeostasis in such patients.

6 **ADVERSE REACTIONS**

6.1 Clinical Trial Experience

Adult Patients

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety data derived from Velphoro clinical trials reflect exposure to Velphoro in 2 active-controlled clinical studies involving a total of 778 adult patients on hemodialysis and 57 adult patients on peritoneal dialysis exposed for up to 55 weeks. Dosage regimens ranged from 250 mg to 3,000 mg per day.

In a parallel design, dose-finding study of Velphoro with a treatment duration of 6 weeks in hemodialysis patients, adverse reactions for Velphoro (N=128) were similar to those reported for the active-control group (sevelamer hydrochicide) (N=26), with the exception of discolored feces (12%) which did not occur in the active-control group. Diarrhea was reported in 6% of patients treated with Velphoro. In a 55-week, open-label, active-controlled, parallel design, safety and efficacy study involving 968 hemodialysis patients and 86 peritoneal dialysis patients treated with either Velphoro (N=707 including 57 peritoneal dialysis patients) or the active-control (sevelamer carbonate) (M=348 including 29 peritoneal dialysis patients), adverse reactions occurring in more than 5% in the Velphoro group were diarrhea (24%), discolored feces (16%), and nausea (10%). The majority of diarrhea events in the Velphoro group were mild and transient,

occurring soon after initiation of treatment, and resolving with continued treatment. Adverse reactions occurred at similar rates in hemodialysis and peritoneal dialysis patients. The most common adverse reactions (>1%) leading to withdrawal were diarrhea (4%), product taste abnormal (2%), and nausea (2%).

Pediatric Patients

In an open-label, randomized study, with a 10-week dose titration period and 24-week safety extension, 60 patients 6 to 18 years of age received at least one dose of Velphoro, including 30 patients (50%) exposed for at least 19 weeks *[see Clinical Studies (14.3)]*. The safety profile of Velphoro in pediatric patients was similar to that observed in adult patients. Velphoro is not approved in pediatric patients 6 years to less than 9 years of age because of the lack of an appropriate dosage strength.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of Velphoro that are not included in other sections of labeling. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. *Gastrointestinal Disorders: tooth discoloration Skin and Subcutaneous Tissue Disorders: rash*

7 DRUG INTERACTIONS

Table 1 Oral drugs that can be add	ninistered concomitantly with Velphoro	
Ciproflovacin		
Digovin		
Englanril		
Eurocomido		
HMC CoA reductore inhibitore		
Hydrochlorothiazido		
Hyulocillololillaziue		
LUSallali		
Nifodinino		
Omonrazolo		
Quinidino		
Warfarin		
Oral drugs that are to be separated from Velphoro		
	Dosing Recommendations	
Doxycycline		
Acetylsalicylic acid	Take at least 1 hour before Velphoro.	
Cephalexin		

Oral medications not listed in Table 1

There are no empirical data on avoiding drug interactions between Velphoro and most concomitant oral drugs. For oral medications where a reduction in the bioavailability of that medication would have a clinically significant effect on its safety or efficacy, consider separating the administration of the two drugs. The necessary separation depends upon the absorption characteristics of the medication concomitantly administered, such as the time to reach peak systemic levels and whether the drug is an immediate-release or an extended-release product. Where possible, consider monitoring for clinical response and/or blood levels of concomitant medications that have a narrow therapeutic range.

Take at least 4 hours before Velphoro.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Levothyroxine

Velphoro is not systemically absorbed following oral administration and maternal use is not expected to result in fetal exposure to the drug.

<u>Data</u> Animal Data

In pregnant rats given up to 800 mg/kg/day Velphoro by oral gavage from Days 6 to 17 post-mating, no embryo-fetal development toxicity was observed. This dose corresponds to 16 times the maximum recommended clinical dose.

In pregnant rabbits given 50, 100 or 200 mg/kg/day Velphoro by oral gavage, from Days 6 to 19 post-mating, the number of fetuses with incomplete/unossified epiphyses and metacarpals/phalanges was increased at the highest dose (corresponding to 4 times the recommended maximum clinical dose). Litter parameters were not adversely affected.

In pregnant rats given Velphoro at 100, 280, or 800 mg/kg/day by oral gavage from Day 6 post-mating to lactation Day 20, offspring body weight gain was lower at age 5–13 weeks and neuromuscular function was delayed at the dose of 800 mg/kg/day. This dose represented 16 times the maximum recommended clinical dose.

8.2 Lactation

Velphoro is not absorbed systemically following oral administration and breastfeeding is not expected to result in exposure of the child to Velphoro.

8.4 Pediatric Use

Velphoro is indicated for the control of serum phosphorus levels in pediatric patients 9 years of age and older with CKD on dialysis. Use of Velphoro for this indication is supported by evidence from an adequate and well-controlled study in adults, with additional pharmacodynamic and safety data in pediatric patients with advanced CKD (an estimated glomerular filtration rate <30 mL/min/1.73 m² or CKD on dialysis) [see Adverse Reactions (6.1) and Clinical Studies (14.3)]. The safety and effectiveness of Velphoro have been established for the control of serum phosphorus levels in pediatric patients 6 years to less than 9 years of age because of the lack of an approved in pediatric patients 6 years to less than 9 years of age because of the lack of an appropriate dosage strength.

The safety and effectiveness of Velphoro have not been established in pediatric patients younger than 6 years of age. Although the study included six patients 2 to <6 years of age who received Velphoro, based on the available data, it is unclear whether the dosing regimen that was evaluated is effective in reducing serum phosphorus in this age group. Velphoro has not been studied in pediatric patients below 2 years of age.

8.5 Geriatric Use

Of the total number of subjects in two active-controlled clinical studies of Velphoro (N=835), 29.7% (n=248) were 65 and over, while 8.7% (n=73) were aged 75 years and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects.

10 OVERDOSAGE

There are no reports of overdosage with Velphoro in patients. Since the absorption of iron from Velphoro is low *[see Clinical Pharmacology (12.3)]*, the risk of systemic iron toxicity is low. Hypophosphatemia should be treated by standard clinical practice.

Velphoro has been studied in doses up to 3,000 mg per day.

11 DESCRIPTION

The Velphoro drug substance is a mixture of polynuclear iron(III)-oxyhydroxide, sucrose, and starches. The active moiety of Velphoro, polynuclear iron(III)-oxyhydroxide, is practically insoluble and cannot be absorbed.

Velphoro chewable tablets for oral use are brown, circular, bi-planar, and are embossed with "PA 500" on 1 side. Each tablet of Velphoro contains 500 mg iron (in 2,500 mg sucroferric oxyhydroxide). One tablet is equivalent to approximately 1.4 g of carbohydrates (750 mg sucrose and 700 mg starches as potato starch and pregelatinized maize starch). The inactive ingredients are berry flavor, neohesperidin dihydrochalcone, magnesium stearate, and silica (colloidal, anhydrous).

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

In the aqueous environment of the GI tract, phosphate binding takes place by ligand exchange between hydroxyl groups and/or water in sucroferric oxyhydroxide and the phosphate in the diet. The bound phosphate is eliminated with feces. Both serum phosphorus levels and calcium-phosphorus product levels are reduced as a consequence of the reduced dietary phosphate absorption.

12.2 Pharmacodynamics

In vitro studies have demonstrated a robust phosphate binding capacity of Velphoro over the physiologically relevant pH range of the GI tract (1.2-7.5). The phosphate binding capacity of Velphoro peaked at pH 2.5, resulting in 96% of the available phosphate being adsorbed (phosphorus:iron concentration ratio 0.4:1).

12.3 Pharmacokinetics

The active moiety of Velphoro, polynuclear iron(III)-oxyhydroxide (pn-Fe00H), is practically insoluble in water and therefore not absorbed and not metabolized. Its degradation product, mononuclear iron species, can however be released from the surface of pn-Fe00H and be absorbed.

Because of the insolubility and degradation characteristics of Velphoro, no classical pharmacokinetic studies can be carried out.

The sucrose and starch components of Velphoro can be digested to glucose and fructose, and maltose and glucose, respectively. These compounds can be absorbed in the blood.

The iron uptake from radiolabeled Velphoro drug substance, 2,000 mg in 1 day, was investigated in 16 chronic kidney disease patients (8 pre-dialysis and 8 hemodialysis patients) and 8 healthy volunteers with low iron stores (serum ferritin <100 mcg/L). In healthy subjects, the median uptake of radiolabeled iron in the blood was 0.43% on Day 21. In chronic kidney disease patients, the median uptake was much less, 0.04% on Day 21.

Drug Interaction Studies

In vitro

In vitro interactions were studied in aqueous solutions which mimic the physicochemical conditions of the gastrointestinal tract with or without the presence of phosphate (400 mg). The study was conducted at pH 3.0, 5.5 and 8.0 with incubation at 37° C for 6 hours.

Interaction with Velphoro was seen with the following drugs: alendronate, doxycycline, acetylsalicylic acid, cephalexin, levothyroxine, and paricalcitol.

The following drugs did not show interaction with Velphoro: ciprofloxacin, enalapril, hydrochlorothiazide, metoprolol, nifedipine, and quinidine.

In vivo

Five *in vivo* drug interaction studies (N = 40/study) were conducted with losartan, furosemide, digoxin, omeprazole and warfarin in healthy subjects receiving 1,000 mg Velphoro 3 times a day with meals. Velphoro did not alter the systemic exposure as measured by the area under the curve (AUC) of the tested drugs when co-administered with Velphoro or given 2 hours later.

Data from the clinical studies (Study-05A and Study-05B) show that Velphoro does not affect the lipid lowering effects of HMG-CoA reductase inhibitors or the PTH lowering effect of calcitriol.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity studies were performed in mice and rats.

In the 2-year carcinogenicity study in mice, animals were given Velphoro by diet at doses of 250, 500 or 1,000 mg/kg/day. There was no clear evidence of a carcinogenic effect in mice. Mucosal hyperplasia, with diverticulum/cyst formation was observed in the colon and caecum of mice after 2 years treatment. In the 2-year rat carcinogenicity study, animals were given Velphoro by diet at doses of 40, 150 or 500 mg/kg/day. No

statistically significantly increased incidences of tumors were found, but there were increased incidences in epithelial hyperplasia with or without submucosal inflammation in duodenum, cecum and colon at the dose of 500 mg/kg/day (10 times the maximum recommended clinical dose).

Velphoro was not mutagenic, clastogenic or DNA damaging *in vitro* in the Ames bacterial reverse mutation test, or in the Chinese-hamster fibroblast chromosomal aberration test, or *in vivo* in the rat Comet assay or peripheral blood micronucleus test.

In rats, mating performance and fertility were unaffected by Velphoro at oral doses up to 800 mg/kg/day (16 times the maximum recommended clinical dose).

14 CLINICAL STUDIES

14.1 Fixed-dose Study

In Study-03A, 154 ESRD adult patients on hemodialysis who were hyperphosphatemic (serum phosphorus >5.5 mg/dL but <7.75 mg/dL) following a 2-week phosphate binder washout period, were randomized to receive Velphoro at 250 mg/day, 1,000 mg/day, 1,500 mg/day, 2,000 mg/day, or 2,500 mg/day or active-control (sevelamer hydrochloride). Velphoro treatment was divided across meals, depending on dose. No dose titration was allowed. Within each of the groups, the serum phosphorus level at the end of treatment was compared to baseline value. Velphoro was shown to be efficacious (p=0.016) for all doses except 250 mg/day. There were no patient-reported dose limiting treatment-emergent adverse events.

Mean changes in iron parameters (ferritin, transferrin saturation (TSAT) and transferrin) and vitamins (A, D, E and K) were generally not clinically meaningful and showed no apparent trends across the treatment groups. Velphoro had a similar GI adverse event profile *[see Adverse Reactions (6.1)]* to sevelamer hydrochloride, and no dose-dependent trend in GI events was observed.

14.2 Dose Titration Study

In Study-05A, 1,055 adult patients on hemodialysis (N=968) or peritoneal dialysis (N=87) with serum phosphorus ≥ 6 mg/dL following a 2-4 week phosphate binder washout period, were randomized and treated with either Velphoro, at a starting dose of 1,000 mg/day (N=707), or active-control (sevelamer carbonate, N=348) for 24 weeks. At the end of Week 24, 93 patients on hemodialysis whose serum phosphorus levels were controlled (<5.5 mg/dL) with Velphoro in the first part of the study, were re-randomized to continue treatment with either their Week 24 maintenance dose (N=44 or a non-effective low dose control 250 mg/day, N=49) of Velphoro for a further 3 weeks. At Week 27, a superiority analysis of the Velphoro maintenance dose versus low dose was performed. The maximum dose of Velphoro was 3,000 mg/day (2 tablets/day) and the minimum dose was 1,000 mg/day (2 tablets/day). Velphoro was administered with food and the daily dose was divided across the largest meals of the day.

The Velphoro maintenance dose (1,000 to 3,000 mg/day) was statistically significantly superior in sustaining the phosphorus lowering effect in hemodialysis patients at Week 27 (p<0.001) compared with the non-effective low dose control. The results are provided in Table 2.

Table 2 Mean (SD) Serum Phosphorus and Change from Baseline to End of Treatment

	Mean (SD) Serum Phosphorus (mg/dL)		
	Velphoro Maintenance Dose (1,000 to 3,000 mg/day) (N = 44)	Velphoro Low Dose Control (250 mg/day) (N = 49)	
Week 24 (BL)	4.7 (1.03)	5.0 (1.14)	
Week 25	4.7 (0.91)	6.3 (1.44)	
Week 26	4.7 (1.21)	6.6 (1.91)	
Week 27/End of Treatment	5.0 (1.07)	6.8 (1.63)	
Change from BL to End of Treatment	0.3 (1.22)*	1.8 (1.47)	

* p < 0.001 for the difference in least square means of the change from BL to Week 27/ End of Treatment (LOCF principle) between Velphoro maintenance dose and low dose using a covariance analysis (MIXED Model).

Notes: BL is Week 24 or latest value available before Week 24 when Week 24 result is missing; End of Treatment is Week 27 value or includes the latest evaluable measurement after Week 24 (i.e., LOCF).

BL = Baseline; LOCF = Last observation carried forward; SD = Standard

deviation.

Following completion of Study-05A, 658 patients (597 on hemodialysis and 61 on peritoneal dialysis) were treated in the 28-week extension study (Study-05B) with either Velphoro (N = 391) or sevelamer carbonate (N = 267) according to their original randomization.

Serum phosphorus levels declined rapidly during the first few weeks of treatment and remained relatively constant thereafter. The phosphorus lowering effect of Velphoro was consistently maintained and controlled through 12 months of treatment (shown in Figure 1). The proportion of adherent patients for Velphoro was 86% at 52 weeks.



Figure 1

Mean change (\pm SEM) from baseline in serum phosphorus over time in Study-05A and extension Study-05B. Insert showing the mean change (\pm SEM) from baseline in serum phosphorus during the withdrawal phase of the study (Weeks 24 to 27) for Velphoro non-effective low dose control (250 mg/day) versus Velphoro maintenance dose.

Age, gender, race, or dialysis modality did not affect the efficacy of Velphoro.

There were no clinically meaningful changes for serum iron, ferritin, TSAT levels, vitamins (A, D, E and K) with Velphoro. There was no evidence of accumulation of iron during one year treatment.

14.3 Pediatric Study

A clinical study of Velphoro (NCT02688764) was conducted in pediatric patients with hyperphosphatemia and advanced CKD (defined as an estimated glomerular filtration rate <30 mL/min/1.73 m² or CKD on dialysis). Following a washout period of up to 3 weeks, 78 patients 6 to 18 years of age were randomized to Velphoro (N=60) or active control calcium acetate (Phoslyra) (N=18) for a 10-week dose titration period, followed by a 24-week safety extension. Approximately 81% of patients were CKD patients on dialysis (69% on hemodialysis and 12% on peritoneal dialysis). 43 patients randomized to Velphoro completed the dose titration period.

The least squares (LS) mean reduction in serum phosphorus levels from baseline to the end of the dose titration period in the Velphoro group (N=59) was -0.52 mg/dL (95% Cl: -0.97, -0.07). The percentage of patients with serum phosphorus levels within normal ranges increased from 39% at baseline to 64% at the end of the dose titration period.

Velphoro is not approved in pediatric patients 6 years to less than 9 years of age because of the lack of an appropriate dosage strength.

The median prescribed daily dose of Velphoro was approximately 1500 mg during the dose titration period for patients 9 to 18 years of age.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Velphoro chewable tablets are brown, circular, bi-planar tablets, embossed with "PA 500" on 1 side. Each tablet of Velphoro contains 500 mg iron as sucroferric oxyhydroxide. Velphoro tablets are packaged as follows: NDC 49230-645-51 Bottle of 90 tablets

16.2 Storage

Keep the bottle tightly closed in order to protect from moisture.

Store at 25°C (77°F) with excursions permitted to 15°C to 30°C (59°F to 86°F). *[See USP Controlled Room Temperature]*

17 PATIENT COUNSELING INFORMATION

Inform patients that Velphoro tablets should be chewed or crushed. Do not swallow whole [see Dosage and Administration (2.2)].

Advise patients that Velphoro should be taken with meals.

Instruct patients on concomitant medications that should be dosed apart from Velphoro *[see Drug Interactions (7)]*. Inform patients that Velphoro can cause discolored (black) stool, but this discoloration of the stool is considered normal with oral medications containing iron. Gastrointestinal bleeding may be masked by stool discoloration.

Inform patients that Velphoro can stain teeth.

Inform patients to report any rash to their health care professional.

Distributed by: Fresenius Medical Care North America 920 Winter Street Waltham, MA 02451 Patents Apply – visit www.fmcna.com/patents © 2014 – 2020, 2024 Fresenius Medical Care North America. All Rights Reserved.